

International Journal of Modern Physics A
 © World Scientific Publishing Company

QUARK-DIQUARK SYSTEMATICS OF BARYONS AND THE $SU(6)$ SYMMETRY FOR LIGHT STATES

A.V. ANISOVICH, V.V. ANISOVICH*, M.A. MATVEEV, V.A. NIKONOV, A.V. SARANTSEV
 and T.O. VULFS

Petersburg Nuclear Physics Institute, Gatchina, 188300 Russia

Received Day Month Year

Revised Day Month Year

We continue our attempts to systematize baryons, which are composed of light quarks ($q = u, d$), as quark-diquark systems. The notion of two diquarks is used: (i) D_1^1 , with the spin $S_D = 1$ and isospin $I_D = 1$ and (ii) D_0^0 , with $S_D = 0$ and $I_D = 0$. Here we try to resolve the problem of the low-lying $\Delta(\frac{5}{2}^-)$ states: in the last experiments the lightest state is observed at $\gtrsim 2200$ MeV, not at $1900 - 2000$ MeV as it has been stated 20 years ago. We are looking for different systematization variants with the forbidden low-lying $\Delta(\frac{5}{2}^-)$ states in the mass region $\lesssim 2000$ MeV. We see that the inclusion of the $SU(6)$ constraints on qD_1^1 states with angular momentum $L = 1$ results in a shift of the lightest $\Delta(\frac{5}{2}^-)$ isobar to ~ 2300 MeV. The scheme with the $SU(6)$ constraints for low-lying qD_1^1 and qD_0^0 states (with $L = 0, 1$) is presented in detail here.

PACS numbers: 11.25.Hf, 123.1K

1. Introduction

Baryons (we mean standard non-exotic baryons) are composite systems of three constituent quarks, each of them in their turn being a complicated system of quarks and gluons. We do not know the detailed structure of baryons, even in the language of constituent quarks, except for a fragmentary knowledge on low-lying states.

We know that low-lying baryons satisfy the $SU(6)$ symmetry, but as to heavier ones we are not certain about. Numerous model calculations, which describe rather well the low-lying states, are at variance in their predictions concerning highly excited states, e.g., see ^{1,2,3,4,5} and references therein. In addition, the number of highly excited states predicted by such models exceeds considerably the number of observed states. One might believe that experimental investigations of baryon spectra were not complete. Nevertheless, it does not remove the question whether it is possible to construct a more effective scheme for highly excited states, with less degrees of freedom and less number of highly excited states. The introduction of another effective particle, that is the diquark, provides us with such a possibility.

*anisovic@thd.pnpi.spb.ru

The size of diquark as a composite quark–antiquark system is believed to be of the order of that of quark, $\sim 0.2\text{--}0.3\text{ fm}$ ^{6,7}. So, it is doubtful if we can interpret the low-lying baryon (a compact system, $\sim 0.7\text{--}0.9\text{ fm}$) as a system of effective diquark and constituent quark.

The idea of baryon as spatially separated quark and diquark can be for sure tried on the highly excited baryon systems which are of a larger size – such an idea was suggested in⁸. But in⁸ we accepted also that the quark–diquark structure does not work for low-lying baryons with angular momentum $L = 0$ – for these states the $SU(6)$ symmetry was applied.

Actually, in⁸ we have made an attempt to systematize highly excited baryons, assuming that they do not like to be formed as three-particle colour quark systems but prefer to be created as two-particle, quark–diquark, compound states: $q_\alpha D^\alpha \equiv q_\alpha [\varepsilon^{\alpha\beta\gamma} q_{\beta\gamma}]$ where $\varepsilon^{\alpha\beta\gamma}$ is a totally antisymmetrical tensor in the colour space.

In⁸ we have considered two schemes of the quark–diquark construction for the $L \geq 1$ states, namely:

- (1) The diquark masses $M_{D_0^0} \neq M_{D_1^1}$.
- (2) The diquark masses are equal to each other $M_{D_0^0} = M_{D_1^1}$; besides, the states qD_0^0 and qD_1^1 with the total spin $S = 1/2$ (here $\mathbf{S} = \mathbf{S}_q + \mathbf{S}_D$) overlap. This scheme decreases essentially the number of highly excited states.

In the quark–diquark scheme of⁸ we face a specific transition to the $SU(6)$ limit. This procedure is in fact a projection of the qD_0^0 and qD_1^1 wave functions on the $SU(6)$ basis – the result depends on the hypothesis on diquark masses. In the $L = 0$ states, at $M_{D_0^0} = M_{D_1^1}$, we have two basic ($n = 1$) states, $N_{\frac{1}{2}^+}(940)$ and $\Delta_{\frac{3}{2}^+}(1240)$, see equation (46) in⁸, while at $M_{D_0^0} \neq M_{D_1^1}$ the additional basic state, $N_{\frac{1}{2}^+}(1440)$, appears (see equation (49) in⁸).

In the present paper we accept the equality of masses of scalar and axial–vector diquarks, $M_{D_0^0} = M_{D_1^1}$. In recent analyses of spectra near the Roper resonance one may find arguments in favour of the existence of one pole in the partial wave P_{11} near 1400 MeV. Analytical properties of the partial wave P_{11} in the best fit of^{9,10} are shown in Fig. 1; the fitting to two poles near 1400 MeV gives us a worse description of data. Of course, the Roper pole splits, owing to the momentum-dependence of width near the $\pi\Delta$ branching point. This splitting is quite similar to that observed in the Flatté formula¹¹ for $f_0(980)$ near the $K\bar{K}$ threshold. Still, we attribute such “satellite poles” to the main one, in case in question, to the Roper pole ($1370 \pm i96$); note that in Fig. 1 the satellite pole is hidden under the $\pi\Delta$ cut.

There is a question to what low-lying states the $SU(6)$ symmetry may be applied and where we have the region with spatially separated quark and diquark. Only the experiment can answer this question. If we turn to the PDG compilation¹², it may seem that the $SU(6)$ symmetry can be applied to $L = 0$ only, while at $L \geq 1$ the domain of quark–diquark structures begin. However, the latest analyses^{9,10,13} give rise to doubts. The matter is that in the experiments carried out in the eighties^{14,15,16} the resonance $\Delta_{5/2^-}(1930 \pm 50)$ has been observed rather definitely.




Fig. 1. Complex- M plane: position of poles of resonance states $N(1440)$ and $N(1710)$. The cuts related to the threshold singularities πN , $\pi\Delta$, ρN and σN are shown by vertical solid lines. The Roper pole is located near the $\pi\Delta$ cut, while its satellite pole is hidden under the cut.

Still, modern analyses^{9,10,13} point to the lower $\Delta_{5/2-}$ state, being located around 2200 MeV or even higher. It gives arguments for expanding the $SU(6)$ symmetry constraints on the $L = 1$ states.

In the present paper we accept that both sets of states ($L = 0$ and $L = 1$) satisfy the $SU(6)$ symmetry requirements.

The paper is organized as follows:

In Section 2 we present wave functions of quark-diquark states and demonstrate how the imposing of the $SU(6)$ constraints affect these systems. In Section 3 we suggest the setting of the $L = 0$ and $L = 1$ states, while states with $L > 1$ are discussed in Section 4. Here we demonstrate the setting of all quark-diquark states on the (J, M^2) and (n, M^2) trajectories.

2. Wave functions of quark-diquark systems and the $SU(6)$ constraints

We present here the diquark wave function, give general form for the quark-diquark wave function and present the scheme of projecting them on the $SU(6)$ basis.

2.1. S -wave diquarks and baryons

Recall that we use two S -wave diquarks with color numbers $\bar{\mathbf{c}} = 3$: scalar diquark D_0^0 and axial-vector one, $D_{1SZ}^{1I_Z}$. The diquark spin-flavor wave functions with

4 *A.V. Anisovich et al.*

$I_D = 1$, $S_D = 1$ and with $I_D = 0$, $S_D = 0$ read as follows:

$$\begin{aligned}
D_{11}^{11}(ij) &= u^\uparrow(i)u^\uparrow(j), \\
D_{10}^{11}(ij) &= \frac{1}{\sqrt{2}} \left(u^\uparrow(i)u^\downarrow(j) + u^\downarrow(i)u^\uparrow(j) \right), \\
D_{11}^{10}(ij) &= \frac{1}{\sqrt{2}} \left(u^\uparrow(i)d^\uparrow(j) + d^\uparrow(i)u^\uparrow(j) \right), \\
D_{10}^{10}(ij) &= \frac{1}{2} \left(u^\uparrow(i)d^\downarrow(j) + u^\downarrow(i)d^\uparrow(j) + d^\uparrow(i)u^\downarrow(j) + d^\downarrow(i)u^\uparrow(j) \right), \\
D_0^0(ij) &= \frac{1}{2} \left(u^\uparrow(i)d^\downarrow(j) - u^\downarrow(i)d^\uparrow(j) - d^\uparrow(i)u^\downarrow(j) + d^\downarrow(i)u^\uparrow(j) \right). \quad (1)
\end{aligned}$$

Let us consider, first, the Δ isobar at $I_Z = 3/2$ with fixed J, J_Z , total spin S and orbital momentum L . The wave function for this state at arbitrary n reads

$$\begin{aligned}
&\sum_{S_Z, m_z} C_{L J_Z - S_Z}^J C_{S S_Z}^S C_{1 S_Z - m_z}^{S S_Z} C_{\frac{1}{2} m_z}^{\frac{3}{2} \frac{3}{2}} C_{\frac{1}{2} \frac{1}{2}}^{\frac{3}{2} \frac{3}{2}} \left(u^{m_z}(1) D_{1 S_Z - m_z}^{11}(23) \right. \\
&\quad \left. \times |\vec{k}_{1 cm}|^L Y_L^{J_Z - S_Z}(\theta_1, \phi_1) \Phi_1^{(L)}(1; 23) + (1 \rightleftharpoons 2) + (1 \rightleftharpoons 3) \right). \quad (2)
\end{aligned}$$

Here $|\vec{k}_{1 cm}|$ and (θ_1, ϕ_1) are the momenta and momentum angles of the first quark in the c.m. system.

For other I_Z , one should include into the wave function a summing over isotopic states which means the following substitution in (2):

$$C_{\frac{1}{2} \frac{1}{2}}^{\frac{3}{2} \frac{3}{2}} u^{m_z}(1) D_{1 S_Z - m_z}^{11}(23) \rightarrow \sum_{j_z} C_{1 I_Z - j_z}^{\frac{3}{2} I_Z} C_{\frac{1}{2} j_z}^{\frac{3}{2} I_Z} q_{j_z}^{m_z}(1) D_{1 S_Z - m_z}^{1 I_Z - j_z}(23). \quad (3)$$

To project the wave functions (2) or (3) on the $SU(6)$ wave function set, we should use symmetrical coordinate/momentum wave functions

$$\Phi_{I_D}^{(L)}(\ell; ij) \rightarrow \Phi_{I_D}^{(L)}(\ell, i, j) = \Phi_{I_D}^{(L)}(j, \ell, i) = \Phi_{I_D}^{(L)}(i, j, \ell). \quad (4)$$

For example, one may accept that $\Phi_{I_D}^{(L)}(\ell, i, j)$ depends on s only, $\Phi_{I_D}^{(L)}(\ell, i, j) \rightarrow \varphi_{I_D}^{(L)}(s)$. Indeed, in this limit we have

$$\begin{aligned}
&\sum_{S_Z, m_z} C_{L J_Z - S_Z}^J C_{S S_Z}^S C_{1 S_Z - m_z}^{S S_Z} C_{\frac{1}{2} m_z}^{\frac{3}{2} I_Z} \sum_{j_z} C_{1 I_Z - j_z}^{\frac{3}{2} I_Z} C_{\frac{1}{2} j_z}^{\frac{3}{2} I_Z} \left(q_{j_z}^{m_z}(1) D_{1 S_Z - m_z}^{1 I_Z - j_z}(23) |\vec{k}_{1 cm}|^L Y_L^{J_Z - S_Z}(\theta_1, \phi_1) \right. \\
&\quad \left. + (1 \rightleftharpoons 2) + (1 \rightleftharpoons 3) \right) \varphi_1^{(L)}(s). \quad (5)
\end{aligned}$$

For nucleon states ($I = 1/2$) we write

$$\begin{aligned}
&\sum_{S_Z, m_z} C_{L J_Z - S_Z}^J C_{S S_Z}^S C_{1 S_Z - m_z}^{S S_Z} C_{\frac{1}{2} m_z}^{\frac{1}{2} I_Z} \sum_{j_z} C_{1 I_Z - j_z}^{\frac{1}{2} I_Z} C_{\frac{1}{2} j_z}^{\frac{1}{2} I_Z} \left(q_{j_z}^{m_z}(1) D_{1 S_Z - m_z}^{1 I_Z - j_z}(23) \right. \\
&\quad \left. \times |\vec{k}_{1 cm}|^L Y_L^{J_Z - S_Z}(\theta_1, \phi_1) \Phi_1^{(L)}(1; 23) + (1 \rightleftharpoons 2) + (1 \rightleftharpoons 3) \right). \quad (6)
\end{aligned}$$

The $SU(6)$ limit, as previously, is reached at $\Phi_1^{(L)}(i; j\ell) \rightarrow \varphi_1^{(L)}(s)$. Then, instead of (6), one has

$$\sum_{S_Z, m_z} C_{L J_Z - S_Z}^{J J_Z} C_{S S_Z}^{S S_Z} C_{1 S_Z - m_z}^{S S_Z} \sum_{j_z} C_{1 I_Z - j_z}^{\frac{1}{2} I_Z} \frac{1}{2} m_z \times \left(q_{j_z}^{m_z}(1) D_{1 S_Z - m_z}^{1 I_Z - j_z}(23) |\vec{k}_{1 cm}|^L Y_L^{J_Z - S_Z}(\theta_1, \phi_1) + (1 \Rightarrow 2) + (1 \Rightarrow 3) \right) \varphi_1^{(L)}(s). \quad (7)$$

For qD_0^0 states the wave function reads in general case as follows:

$$\sum_{m_z} C_{L J_Z - m_z}^{J J_Z} \frac{1}{2} m_z \left(q_{I_z}^{m_z}(1) D_0^0(23) |\vec{k}_{1 cm}|^L Y_L^{J_Z - m_z}(\theta_1, \phi_1) \Phi_0^{(L)}(1; 23) + (1 \Rightarrow 2) + (1 \Rightarrow 3) \right). \quad (8)$$

In the $SU(6)$ limit we have

$$\sum_{m_z} C_{L J_Z - m_z}^{J J_Z} \frac{1}{2} m_z \left(q_{I_z}^{m_z}(1) D_0^0(23) |\vec{k}_{1 cm}|^L Y_L^{J_Z - m_z}(\theta_1, \phi_1) + (1 \Rightarrow 2) + (1 \Rightarrow 3) \right) \varphi_0^{(L)}(s). \quad (9)$$

Baryons are characterized by I and J^P , these states with different S and L and fixed I and J^P can mix. To select independent states, one may orthogonalize wave functions with the same isospin and J^P . The orthogonalization depends on the structure of the momentum/coordinate parts $\Phi_1^{(L)}(i; j\ell)$. But in case of the $SU(6)$ limit the momentum/coordinate wave functions transform in the common factor $\Phi_0^{(L)}(i; j\ell) \rightarrow \varphi_0^{(L)}(i, j, \ell)$ or $\Phi_1^{(L)}(i; j\ell) \rightarrow \varphi_{SU(6)}^{(L)}(s)$, and one should orthogonalize the spin/momentum factors. Namely, let us denote the $SU(6)$ spin/momentum factor of the wave function as $Q_{J^P}^{(A)}$. Then

$$\Psi_{J^P}^{(A)} = Q_{J^P}^{(A)} \varphi_{SU(6)}^{(A)}(s), \quad (10)$$

where $A = I, II, III, \dots$ belong to different (S, L) . The orthogonal set of operators $Q_{J^P}^{(A)}$ is constructed in a standard way:

$$\begin{aligned} Q_{J^P}^{(\perp I)} &\equiv Q_{J^P}^{(I)}, \\ Q_{J^P}^{(\perp II)} &= Q_{J^P}^{(II)} - Q_{J^P}^{(\perp I)} \frac{\left(Q_{J^P}^{(\perp I)} + Q_{J^P}^{(II)} \right)}{\left(Q_{J^P}^{(\perp I)} + Q_{J^P}^{(\perp I)} \right)}, \\ Q_{J^P}^{(\perp III)} &= Q_{J^P}^{(III)} - Q_{J^P}^{(\perp I)} \frac{\left(Q_{J^P}^{(\perp I)} + Q_{J^P}^{(III)} \right)}{\left(Q_{J^P}^{(\perp I)} + Q_{J^P}^{(\perp I)} \right)} - Q_{J^P}^{(\perp II)} \frac{\left(Q_{J^P}^{(\perp II)} + Q_{J^P}^{(III)} \right)}{\left(Q_{J^P}^{(\perp II)} + Q_{J^P}^{(\perp II)} \right)}, \end{aligned} \quad (11)$$

and so on. The convolution of operators $\left(Q_{J^P}^{(A)} + Q_{J^P}^{(B)} \right)$ includes the summation over quark spins as well as integration over quark momenta.

2.2. Wave functions in the $SU(6)$ limit

We present wave functions of the qD_0^0 and qD_1^1 systems in the $SU(6)$ limit, assuming $M_{D_0^0}^2 = M_{D_1^1}^2 \equiv M^2$. The transition $M_{D_0^0}^2 \rightarrow M_{D_1^1}^2$ gives us additional reduction of number of states.

Recall that in the $SU(6)$ limit the momentum/coordinate factors of wave functions, for example, such as $\Phi_0^{(L)}(3, 12) = \Phi_0^{(L)}(s, s_{12})$ and $\Phi_1^{(L)}(3, 12) = \Phi_1^{(L)}(s, s_{12})$, transform into s -dependent ones:

$$\begin{aligned}\Phi_0^{(L)}(s, s_{12} \rightarrow M_{D_0^0}^2) &= \varphi_0^{(L)}(s, M_D^2) \equiv \varphi_0^{(L)}(s), \\ \Phi_1^{(L)}(s, s_{12} \rightarrow M_{D_1^1}^2) &= \varphi_1^{(L)}(s, M_D^2) \equiv \varphi_1^{(L)}(s).\end{aligned}\quad (12)$$

For our purpose, that is, for checking the number of non-vanishing states, we calculate below the wave functions with $I = I_z$ and $J = J_z$ only.

3. Wave functions of the ($L = 1$) states in the $SU(6)$ limit and $M_{D_0^0}^2 \rightarrow M_{D_1^1}^2$

In this sector we have only seven basic states, namely,

- (1) two with $I = 1/2$: $N_{\frac{1}{2}-}(D_0^0) = N_{\frac{1}{2}-}(D_1^1; S = 1/2)$,
 $N_{\frac{3}{2}-}(D_0^0) = N_{\frac{3}{2}-}(D_1^1; S = 1/2)$,
- (2) three with $I = 1/2$: $N_{\frac{1}{2}-}(D_1^1; S = 3/2)$, $N_{\frac{3}{2}-}(D_1^1; S = 3/2)$,
 $N_{\frac{5}{2}-}(D_1^1; S = 3/2)$,
- (3) two with $I = 3/2$: $\Delta_{\frac{1}{2}-}(D_1^1; S = 1/2)$, $\Delta_{\frac{3}{2}-}(D_1^1; S = 1/2)$.

Below the wave functions are written in c.m. system using the following notations for quark momenta: $\vec{k}_1 + \vec{k}_2 + \vec{k}_3 = 0$, $k_{a\pm} = (k_{ax} \pm ik_{ay})/\sqrt{2}$.

(1) Two qD_0^0 systems with $I = 1/2$

We have two basic states:

$$\begin{aligned}4\sqrt{\pi} \cdot \Psi_0^{(L=1)}\left(I = \frac{1}{2}, I_z = \frac{1}{2}, J^P = \frac{1}{2}^-, J_z = \frac{1}{2}\right) \\ = \left(u_1^\uparrow u_2^\uparrow d_3^\downarrow k_{3z} + u_1^\uparrow u_2^\downarrow d_3^\uparrow k_{1z} - \sqrt{2}u_1^\uparrow u_2^\downarrow d_3^\downarrow k_{2+} + u_1^\uparrow d_2^\uparrow u_3^\downarrow k_{1z} + u_1^\uparrow d_2^\downarrow u_3^\uparrow k_{2z} \right. \\ \left. - \sqrt{2}u_1^\uparrow d_2^\downarrow u_3^\downarrow k_{3+} + u_1^\downarrow u_2^\uparrow d_3^\uparrow k_{2z} - \sqrt{2}u_1^\downarrow u_2^\uparrow d_3^\downarrow k_{1+} - \sqrt{2}u_1^\downarrow u_2^\downarrow d_3^\uparrow k_{3+} \right. \\ \left. + u_1^\downarrow d_2^\uparrow u_3^\uparrow k_{3z} - \sqrt{2}u_1^\downarrow d_2^\uparrow u_3^\downarrow k_{2+} - \sqrt{2}u_1^\downarrow d_2^\downarrow u_3^\uparrow k_{1+} + d_1^\uparrow u_2^\uparrow u_3^\downarrow k_{2z} + d_1^\uparrow u_2^\downarrow u_3^\uparrow k_{3z} \right. \\ \left. - \sqrt{2}d_1^\uparrow u_2^\downarrow u_3^\downarrow k_{1+} + d_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1z} - \sqrt{2}d_1^\downarrow u_2^\uparrow u_3^\downarrow k_{3+} - \sqrt{2}d_1^\downarrow u_2^\downarrow u_3^\uparrow k_{2+}\right) \varphi_0^{(L=1)}(s) \quad (13)\end{aligned}$$

and

$$\begin{aligned}4\sqrt{\pi} \cdot \Psi_0^{(L=1)}\left(I = \frac{1}{2}, I_z = \frac{1}{2}, J^P = \frac{3}{2}^-, J_z = \frac{3}{2}\right) \\ = \sqrt{3}\left(u_1^\uparrow u_2^\uparrow d_3^\downarrow k_{3+} + u_1^\uparrow u_2^\downarrow d_3^\uparrow k_{1+} + u_1^\uparrow d_2^\uparrow u_3^\downarrow k_{1+} + u_1^\uparrow d_2^\downarrow u_3^\uparrow k_{2+} + u_1^\downarrow u_2^\uparrow d_3^\uparrow k_{2+} \right. \\ \left. + u_1^\downarrow d_2^\uparrow u_3^\uparrow k_{3+} + d_1^\uparrow u_2^\uparrow u_3^\downarrow k_{2+} + d_1^\uparrow u_2^\downarrow u_3^\uparrow k_{3+} + d_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1+}\right) \varphi_0^{(L=1)}(s). \quad (14)\end{aligned}$$

The same angular momentum/spin wave functions can be constructed with use of the qD_1^1 system, in this way we have two nucleons with $S = 1/2$:

$$\begin{aligned}
& 4\sqrt{\pi} \cdot \Psi_1^{(L=1)} \left(I = \frac{1}{2}, I_z = \frac{1}{2}, S = 1/2, J^P = \frac{1}{2}^-, J_z = \frac{1}{2} \right) \\
&= -\sqrt{2} \left(u_1^\uparrow u_2^\uparrow d_3^\downarrow k_{3z} + u_1^\uparrow u_2^\downarrow d_3^\uparrow k_{1z} - \sqrt{2} u_1^\uparrow u_2^\downarrow d_3^\downarrow k_{2+} + u_1^\uparrow d_2^\uparrow u_3^\downarrow k_{1z} + u_1^\uparrow d_2^\downarrow u_3^\uparrow k_{2z} \right. \\
&\quad - \sqrt{2} u_1^\uparrow d_2^\downarrow u_3^\downarrow k_{3+} + u_1^\downarrow u_2^\uparrow d_3^\uparrow k_{2z} - \sqrt{2} u_1^\downarrow u_2^\uparrow d_3^\downarrow k_{1+} - \sqrt{2} u_1^\downarrow u_2^\downarrow d_3^\uparrow k_{3+} + u_1^\downarrow d_2^\uparrow u_3^\uparrow k_{3z} \\
&\quad - \sqrt{2} u_1^\downarrow d_2^\uparrow u_3^\downarrow k_{2+} - \sqrt{2} u_1^\downarrow d_2^\downarrow u_3^\uparrow k_{1+} + d_1^\uparrow u_2^\uparrow u_3^\downarrow k_{2z} + d_1^\uparrow u_2^\downarrow u_3^\uparrow k_{3z} \\
&\quad \left. - \sqrt{2} d_1^\uparrow u_2^\downarrow u_3^\downarrow k_{1+} + d_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1z} - \sqrt{2} d_1^\downarrow u_2^\uparrow u_3^\downarrow k_{3+} - \sqrt{2} d_1^\downarrow u_2^\downarrow u_3^\uparrow k_{2+} \right) \varphi_1^{(L=1)}(s) \quad (15)
\end{aligned}$$

and

$$\begin{aligned}
& 4\sqrt{\pi} \cdot \Psi_1^{(L=1)} \left(I = \frac{1}{2}, I_z = \frac{1}{2}, S = 1/2, J^P = \frac{3}{2}^-, J_z = \frac{3}{2} \right) = \\
&= \sqrt{3} \left(u_1^\uparrow u_2^\uparrow d_3^\downarrow k_{3+} + u_1^\uparrow u_2^\downarrow d_3^\uparrow k_{1+} + u_1^\uparrow d_2^\uparrow u_3^\downarrow k_{1+} + u_1^\uparrow d_2^\downarrow u_3^\uparrow k_{2+} + u_1^\downarrow u_2^\uparrow d_3^\uparrow k_{2+} \right. \\
&\quad \left. + u_1^\downarrow d_2^\uparrow u_3^\uparrow k_{3+} + d_1^\uparrow u_2^\uparrow u_3^\downarrow k_{2+} + d_1^\uparrow u_2^\downarrow u_3^\uparrow k_{3+} + d_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1+} \right) \varphi_1^{(L=1)}(s). \quad (16)
\end{aligned}$$

The wave function of (13) coincides with that of (15), while the wave function of (14) coincides with (16), because we use equation (12): $\varphi_0^{(L=1)}(1, 2, 3) = \varphi_1^{(L=1)}(1, 2, 3)$. In this case we deal with two (not four) states.

(2) Three qD_1^1 systems with $I = 1/2$

There are three states with $S = 3/2$:

$$\begin{aligned}
& 4\sqrt{\pi} \cdot \Psi_1^{(L=1)} \left(I = \frac{1}{2}, I_z = \frac{1}{2}, S = 3/2, J^P = \frac{1}{2}^-, J_z = \frac{1}{2} \right) \\
&= \left(3u_1^\uparrow u_2^\uparrow d_3^\uparrow k_{3-} - \sqrt{2} u_1^\uparrow u_2^\uparrow d_3^\downarrow k_{3z} - \sqrt{2} u_1^\uparrow u_2^\downarrow d_3^\uparrow k_{3z} - u_1^\uparrow u_2^\downarrow d_3^\downarrow k_{3+} + 3u_1^\uparrow d_2^\uparrow u_3^\uparrow k_{2-} \right. \\
&\quad - \sqrt{2} u_1^\uparrow d_2^\downarrow u_3^\downarrow k_{2z} - \sqrt{2} u_1^\downarrow u_2^\uparrow d_3^\uparrow k_{2z} - u_1^\downarrow d_2^\downarrow u_3^\downarrow k_{2+} - \sqrt{2} u_1^\downarrow u_2^\uparrow d_3^\uparrow k_{3z} \\
&\quad - u_1^\downarrow u_2^\downarrow d_3^\downarrow k_{3+} - u_1^\downarrow u_2^\downarrow d_3^\uparrow k_{3+} - \sqrt{2} u_1^\downarrow d_2^\uparrow u_3^\uparrow k_{2z} - u_1^\downarrow d_2^\uparrow u_3^\downarrow k_{2+} \\
&\quad - u_1^\downarrow d_2^\downarrow u_3^\uparrow k_{2+} + 3d_1^\uparrow u_2^\uparrow u_3^\uparrow k_{1-} - \sqrt{2} d_1^\uparrow u_2^\uparrow u_3^\downarrow k_{1z} - \sqrt{2} d_1^\uparrow u_2^\downarrow u_3^\uparrow k_{1z} \\
&\quad \left. - d_1^\uparrow u_2^\downarrow u_3^\downarrow k_{1+} - \sqrt{2} d_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1z} - d_1^\downarrow u_2^\uparrow u_3^\downarrow k_{1+} - d_1^\downarrow u_2^\downarrow u_3^\uparrow k_{1+} \right) \varphi_1^{(L=1)}(s), \quad (17)
\end{aligned}$$

$$\begin{aligned}
& 4\sqrt{\pi} \cdot \Psi_1^{(L=1)} \left(I = \frac{1}{2}, I_z = \frac{1}{2}, S = 3/2, J^P = \frac{3}{2}^-, J_z = \frac{3}{2} \right) \\
&= \frac{3}{\sqrt{15}} \left(-3\sqrt{2} u_1^\uparrow u_2^\uparrow d_3^\uparrow k_{3z} - 2u_1^\uparrow u_2^\uparrow d_3^\downarrow k_{3+} - 2u_1^\uparrow u_2^\downarrow d_3^\uparrow k_{3+} - 3\sqrt{2} u_1^\uparrow d_2^\uparrow u_3^\uparrow k_{2z} \right. \\
&\quad - 2u_1^\uparrow d_2^\downarrow u_3^\downarrow k_{2+} - 2u_1^\downarrow u_2^\uparrow d_3^\uparrow k_{2+} - 2u_1^\downarrow u_2^\uparrow d_3^\downarrow k_{3+} - 2u_1^\downarrow d_2^\uparrow u_3^\uparrow k_{2+} \\
&\quad \left. - 3\sqrt{2} d_1^\uparrow u_2^\uparrow u_3^\uparrow k_{1z} - 2d_1^\uparrow u_2^\uparrow u_3^\downarrow k_{1+} - 2d_1^\uparrow u_2^\downarrow u_3^\uparrow k_{1+} - 2d_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1+} \right) \varphi_1^{(L=1)}(s), \quad (18)
\end{aligned}$$

8 *A.V. Anisovich et al.*

and

$$4\sqrt{\pi} \cdot \Psi_1^{(L=1)} \left(I = \frac{1}{2}, I_z = \frac{1}{2}, S = 3/2, J^P = \frac{5}{2}^-, J_z = \frac{5}{2} \right) \\ = 3\sqrt{2} \left(-u_1^\uparrow u_2^\uparrow d_3^\uparrow k_{3+} - u_1^\uparrow d_2^\uparrow u_3^\uparrow k_{2+} - d_1^\uparrow u_2^\uparrow u_3^\uparrow k_{1+} \right) \varphi_1^{(L=1)}(s). \quad (19)$$

(3) Two qD_1^1 states with $I = 3/2$

In the $(I = \frac{3}{2}, L = 1)$ sector of qD_1^1 states, there are two baryons only in the $SU(6)$ limit – the states with $S = \frac{1}{2}$:

$$4\sqrt{\pi} \cdot \Psi_1^{(L=1)} \left(I = \frac{3}{2}, I_z = \frac{3}{2}, S = 1/2, J^P = \frac{1}{2}^-, J_z = \frac{1}{2} \right) \\ = \left(-\sqrt{6}u_1^\uparrow u_2^\uparrow u_3^\downarrow k_{3z} - \sqrt{6}u_1^\uparrow u_2^\downarrow u_3^\uparrow k_{2z} - \sqrt{6}u_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1z} \right. \\ \left. + 2\sqrt{3}u_1^\uparrow u_2^\downarrow u_3^\downarrow k_{1+} + 2\sqrt{3}u_1^\downarrow u_2^\uparrow u_3^\downarrow k_{2+} + 2\sqrt{3}u_1^\downarrow u_2^\downarrow u_3^\uparrow k_{3+} \right) \varphi_1^{(L=1)}(s), \quad (20)$$

and

$$4\sqrt{\pi} \cdot \Psi_1^{(L=1)} \left(I = \frac{3}{2}, I_z = \frac{3}{2}, S = 1/2, J^P = \frac{3}{2}^-, J_z = \frac{3}{2} \right) \\ = -3\sqrt{2} \left(u_1^\uparrow u_2^\uparrow u_3^\downarrow k_{3+} + u_1^\uparrow u_2^\downarrow u_3^\uparrow k_{2+} + u_1^\downarrow u_2^\uparrow u_3^\uparrow k_{1+} \right) \varphi_1^{(L=1)}(s). \quad (21)$$

3.1. The vanishing class of $(S = 3/2, I = 3/2)$ states

In the $SU(6)$ limit, the states with $S = \frac{3}{2}, I = \frac{3}{2}$ (i.e. $\Delta_{\frac{1}{2}}^-, \Delta_{\frac{3}{2}}^-, \Delta_{\frac{5}{2}}^-$) are absent. It is easy to be certain of that, after calculating wave functions with $J_z = J$. Indeed,

$$\Psi_1^{(L=1)} \left(I = \frac{3}{2}, I_z = \frac{3}{2}, S = 3/2, J^P = \frac{1}{2}^-, J_z = \frac{1}{2} \right) = 0, \\ \Psi_1^{(L=1)} \left(I = \frac{3}{2}, I_z = \frac{3}{2}, S = 3/2, J^P = \frac{3}{2}^-, J_z = \frac{3}{2} \right) = 0, \\ \Psi_1^{(L=1)} \left(I = \frac{3}{2}, I_z = \frac{3}{2}, S = 3/2, J^P = \frac{5}{2}^-, J_z = \frac{5}{2} \right) = 0. \quad (22)$$

The absence of $\Delta_{\frac{5}{2}}^-$, which, being in the $L = 1$ set, should be located at ~ 1900 MeV, is a primary motivation for expanding the $SU(6)$ symmetry onto $L = 1$ states.

4. The $SU(6)$ symmetry and the $L = 0$ and $L = 1$ states

First, we recall the setting of $L = 0$ states, then we discuss the situation with $L = 1$ states under the $SU(6)$ symmetry.

4.1. The $SU(6)$ symmetry for the nucleon $N_{\frac{1}{2}}^+(940)$, isobar

$\Delta_{\frac{3}{2}}^+(1238)$ and their radial excitations

We accept $M_{D_0^1}^2 = M_{D_1^1}^2$ and suppose the $SU(6)$ symmetry for the lowest baryons with $L = 0$. It gives us two ground states – the nucleon $N_{\frac{1}{2}}^+(940)$ and isobar

$\Delta_{\frac{3}{2}+}(1238)$ as well as their radial excitations:

$$\begin{array}{c|cc}
 L=0 & S=\frac{1}{2}, N(\frac{1}{2}^+) & S=\frac{3}{2}, \Delta(\frac{3}{2}^+) \\
 n=1 & 938 \pm 2 & 1232 \pm 4 \\
 n=2 & 1440 \pm 40 & 1635 \pm 75 \\
 n=3 & 1710 \pm 30 & \sim 1880 \\
 n=4 & 2090 \pm 100 & \sim 2150
 \end{array} \quad (23)$$

Note that the mass-squared splitting of the nucleon radial excitation states, $\delta_n M^2(N_{\frac{1}{2}+})$, is of the order of $1.05 \pm 0.15 \text{ GeV}^2$. This value is close to that observed in meson sector^{7,17}:

$$\begin{aligned}
 & M^2[N_{\frac{1}{2}+}(1440)] - M^2[N_{\frac{1}{2}+}(940)] \\
 & \simeq M^2[N_{\frac{1}{2}+}(1710)] - M^2[N_{\frac{1}{2}+}(1440)] \equiv \delta_n M^2(N_{\frac{1}{2}+}) \simeq 1.0 \text{ GeV}^2.
 \end{aligned} \quad (24)$$

The state with $n=4$ cannot be unambiguously determined.

One can see that the mass-squared splitting of $\Delta_{\frac{3}{2}+}$ isobars, $\delta_n M^2(\Delta_{\frac{3}{2}+})$, coincides with that of the nucleon, $\delta_n M^2(N_{\frac{1}{2}+})$, with a good accuracy:

$$\delta_n M^2(\Delta_{\frac{3}{2}+}) = 1.07 \pm 0.05. \quad (25)$$

Let us emphasize that state $\Delta_{\frac{3}{2}+}(1920)$ is classified as $S=3/2, L=2$ states, with $n=1$ (see Section 4). However, this resonance can be reliably classified as radial excitations of $\Delta_{\frac{3}{2}+}(1232)$, with $n=3$. Actually, it means that around $\sim 1920 \text{ MeV}$ one may expect the double-pole structure.

4.2. Setting of the $L=1$ states under the $SU(6)$ symmetry constraints

$$\begin{array}{c|c|c|c|c|c}
 L=1 & S=\frac{1}{2} & \left| \begin{array}{c} n=1 \\ n=2 \\ n=3 \\ n=4 \end{array} \right| & \begin{array}{c} N(\frac{1}{2}^-) \\ (1535 \pm 20) \\ (1905 \pm 60) \\ (2090 \pm 150) \\ \sim 2390 \end{array} & \begin{array}{c} N(\frac{3}{2}^-) \\ (1524 \pm 5) \\ (1870 \pm 25) \\ (2160 \pm 35) \\ \sim 2390 \end{array} & \\
 & S=\frac{3}{2} & \left| \begin{array}{c} n=1 \\ n=2 \\ n=3 \\ n=4 \end{array} \right| & \begin{array}{c} N(\frac{1}{2}^-) \\ (1680 \pm 40) \\ \sim 2010 \\ \sim 2270 \\ \sim 2500 \end{array} & \begin{array}{c} N(\frac{3}{2}^-) \\ (1730 \pm 40) \\ \sim 2000 \\ \sim 2270 \\ \sim 2500 \end{array} & \begin{array}{c} N(\frac{5}{2}^-) \\ (1680 \pm 10) \\ \sim 2000 \\ \sim 2260 \\ \sim 2500 \end{array}
 \end{array} \quad (26)$$

$$\begin{array}{c|c|c|c|c}
 L=1 & S=\frac{1}{2} & \left| \begin{array}{c} n=1 \\ n=2 \\ n=3 \\ n=4 \end{array} \right| & \begin{array}{c} \Delta(\frac{1}{2}^-) \\ (1625 \pm 10) \\ (1910 \pm 50) \\ (2150 \pm 50) \\ \sim 2460 \end{array} & \begin{array}{c} \Delta(\frac{3}{2}^-) \\ (1720 \pm 50) \\ (1995 \pm 40) \\ \sim 2210 \\ \sim 2450 \end{array}
 \end{array} \quad (27)$$

5. The setting of states with $L \geq 2$

For a "naive observer", who does not perform an analysis of double pole structure, the number of states with $S = 1/2$ decreases twice.

In Fig. 2, following ^{9,10,18,12}, we show (J, M^2) plots as they look like for naive observers, while Fig. 3 demonstrates the (J, M^2) plot for ground states ($n = 1$) only.

We have three groups of states with $I = \frac{1}{2}$:

$$\begin{aligned}
 1) \quad & N_{JP} \left(D_0^0; S = \frac{1}{2}, J = L - \frac{1}{2} \right), \quad N_{JP} \left(D_0^0; S = \frac{1}{2}, J = L - \frac{1}{2} \right), \\
 2) \quad & N_{JP} \left(D_1^1; S = \frac{1}{2}, J = L - \frac{1}{2} \right), \quad N_{JP} \left(D_1^1; S = \frac{1}{2}, J = L - \frac{1}{2} \right), \\
 3) \quad & N_{JP} \left(D_1^1; S = \frac{3}{2}, J = L - \frac{3}{2} \right), \quad N_{JP} \left(D_1^1; S = \frac{3}{2}, J = L - \frac{1}{2} \right), \\
 & N_{JP} \left(D_1^1; S = \frac{3}{2}, J = L + \frac{1}{2} \right), \quad N_{JP} \left(D_1^1; S = \frac{3}{2}, J = L + \frac{3}{2} \right),
 \end{aligned} \tag{28}$$

and two with $I = \frac{3}{2}$:

$$\begin{aligned}
 1) \quad & \Delta_{JP} \left(D_1^1; S = \frac{1}{2}, J = L - \frac{1}{2} \right), \quad \Delta_{JP} \left(D_1^1; S = \frac{1}{2}, J = L - \frac{1}{2} \right), \\
 2) \quad & \Delta_{JP} \left(D_1^1; S = \frac{3}{2}, J = L - \frac{3}{2} \right), \quad \Delta_{JP} \left(D_1^1; S = \frac{3}{2}, J = L - \frac{1}{2} \right), \\
 & \Delta_{JP} \left(D_1^1; S = \frac{3}{2}, J = L + \frac{1}{2} \right), \quad \Delta_{JP} \left(D_1^1; S = \frac{3}{2}, J = L + \frac{3}{2} \right),
 \end{aligned} \tag{29}$$

We suppose that in $I = \frac{1}{2}$ sector the states of groups (1) and (2) with the same J and L have the equal masses – resonances are overlapping:

$$\begin{aligned}
 M_{N_{JP}(D_0^0; S=\frac{1}{2}, J=L-\frac{1}{2})} &= M_{N_{JP}(D_1^1; S=\frac{1}{2}, J=L-\frac{1}{2})}, \\
 M_{N_{JP}(D_0^0; S=\frac{1}{2}, J=L+\frac{1}{2})} &= M_{N_{JP}(D_1^1; S=\frac{1}{2}, J=L+\frac{1}{2})}.
 \end{aligned} \tag{30}$$




Fig. 2. Baryon settings on (J^P, M^2) planes in the model with overlapping $qD_0^0(S = 1/2)$ and $qD_1^1(S = 1/2)$ states. Notations are as follows: 1) open squares: predicted $S = 1/2$ ground states, 2) open stars: predicted $S = 3/2$ ground states, 3) open circles: predicted radial excitation states, 4) full triangles: observed states.

12 *A.V. Anisovich et al.*




Fig. 3. Basic baryon settings ($n = 1$) on (J^P, M^2) planes in the model with overlapping $qD_0^0(S = 1/2)$ and $qD_1^1(S = 1/2)$ states (notations are as in Fig 2).

For $P = +$ states the values of baryon masses in the $I = 1/2$ sector are as follows:

$L = 2$	$S = \frac{1}{2}$	$n = 1$		$N(\frac{3}{2}^+)$ (1770 $\pm$ 100)	$N(\frac{5}{2}^+)$ (1685 $\pm$ 5)	
		$n = 2$		(1960 $\pm$ 30)	(2000 $\pm$ 100)	
		$n = 3$		~ 2290	~ 2290	
		$n = 4$		~ 2520	~ 2520	
	$S = \frac{3}{2}$	$n = 1$	$N(\frac{1}{2}^+)$ (1890 $\pm$ 50)	$N(\frac{3}{2}^+)$ ~ 1880	$N(\frac{5}{2}^+)$ ~ 1880	$N(\frac{7}{2}^+)$ (2020 $\pm$ 60)
		$n = 2$	~ 2150	~ 2150	~ 2150	~ 2150
		$n = 3$	~ 2390	~ 2390	~ 2390	~ 2390
		$n = 4$	~ 2610	~ 2610	~ 2610	~ 2610
$L = 4$	$S = \frac{1}{2}$	$n = 1$		$N(\frac{7}{2}^+)$ ~ 2290	$N(\frac{9}{2}^+)$ (2260 $\pm$ 60)	
		$n = 2$		~ 2520	~ 2520	
		$n = 3$		~ 2730	~ 2730	
	$S = \frac{3}{2}$	$n = 1$	$N(\frac{5}{2}^+)$ ~ 2390	$N(\frac{7}{2}^+)$ ~ 2390	$N(\frac{9}{2}^+)$ ~ 2390	$N(\frac{11}{2}^+)$ ~ 2390
		$n = 2$	~ 2610	~ 2610	~ 2610	~ 2610
		$n = 3$	~ 2820	~ 2820	~ 2820	~ 2820
	$S = \frac{1}{2}$	$n = 1$		$N(\frac{11}{2}^+)$ ~ 2730	$N(\frac{13}{2}^+)$ (2700 $\pm$ 100)	
		$n = 2$		~ 2930	~ 2930	
$L = 6$	$S = \frac{1}{2}$	$n = 1$		$N(\frac{9}{2}^+)$ ~ 2820	$N(\frac{11}{2}^+)$ ~ 2820	$N(\frac{13}{2}^+)$ ~ 2820
		$n = 2$	~ 3000	~ 3000	~ 3000	~ 3000
	$S = \frac{3}{2}$	$n = 1$		$N(\frac{11}{2}^+)$ ~ 2730	$N(\frac{13}{2}^+)$ (2700 $\pm$ 100)	
		$n = 2$		~ 2930	~ 2930	
	$S = \frac{1}{2}$	$n = 1$		$N(\frac{9}{2}^+)$ ~ 2820	$N(\frac{11}{2}^+)$ ~ 2820	$N(\frac{13}{2}^+)$ ~ 2820
		$n = 2$	~ 3000	~ 3000	~ 3000	~ 3000
	$S = \frac{3}{2}$	$n = 1$		$N(\frac{11}{2}^+)$ ~ 2730	$N(\frac{13}{2}^+)$ (2700 $\pm$ 100)	
		$n = 2$		~ 2930	~ 2930	

(31)

and for negative-parity states:

$L = 3$	$S = \frac{1}{2}$		$N(\frac{5}{2}^-)$	$N(\frac{7}{2}^-)$		
		$n = 1$	(2160 ± 80)	(2150 ± 30)		
		$n = 2$	~ 2390	~ 2390		
		$n = 3$	~ 2610	~ 2610		
		$n = 4$	~ 2820	~ 2820		
	$S = \frac{3}{2}$		$N(\frac{3}{2}^-)$	$N(\frac{5}{2}^-)$	$N(\frac{7}{2}^-)$	$N(\frac{9}{2}^-)$
		$n = 1$	~ 2260	~ 2260	~ 2260	(2250 ± 50)
		$n = 2$	~ 2490	~ 2490	~ 2490	~ 2490
		$n = 3$	~ 2700	~ 2700	~ 2700	~ 2700
		$n = 4$	~ 2900	~ 2900	~ 2900	~ 2900
$L = 5$	$S = \frac{1}{2}$		$N(\frac{9}{2}^-)$	$N(\frac{11}{2}^-)$		
		$n = 1$	~ 2610	~ 2610		
		$n = 2$	~ 2820	~ 2820		
		$n = 3$	~ 3000	~ 3000		
	$S = \frac{3}{2}$		$N(\frac{7}{2}^-)$	$N(\frac{9}{2}^-)$	$N(\frac{11}{2}^-)$	$N(\frac{13}{2}^-)$
		$n = 1$	~ 2700	~ 2700	~ 2701	~ 2700
		$n = 2$	~ 2900	~ 2900	~ 2898	~ 2900
		$n = 3$	~ 3080	~ 3080	~ 3082	~ 3080

(32)

14 *A.V. Anisovich et al.*In the $I = 3/2$ sector we have for $P = +$ states:

$L = 2$	$S = \frac{1}{2}$	$n = 1$	$\Delta(\frac{3}{2}^+)$	$\Delta(\frac{5}{2}^+)$		
		$n = 2$	~ 1750	~ 1750		
		$n = 3$	~ 2040	~ 2040		
		$n = 4$	~ 2290	~ 2290		
	$S = \frac{3}{2}$	$n = 1$	$\Delta(\frac{1}{2}^+)$	$\Delta(\frac{3}{2}^+)$	$\Delta(\frac{5}{2}^+)$	$\Delta(\frac{7}{2}^+)$
		$n = 2$	(1935 ± 90)	(1995 ± 40)	(1885 ± 25)	(1928 ± 8)
		$n = 3$	~ 2150	~ 2151	~ 2150	~ 2150
		$n = 4$	~ 2400	~ 2400	~ 2400	~ 2400
$L = 4$	$S = \frac{1}{2}$	$n = 1$		$\Delta(\frac{7}{2}^+)$	$\Delta(\frac{9}{2}^+)$	
		$n = 2$		~ 2290	~ 2290	
		$n = 3$		~ 2520	~ 2520	
		$n = 4$		~ 2730	~ 2730	
	$S = \frac{3}{2}$	$n = 1$	$\Delta(\frac{5}{2}^+)$	$\Delta(\frac{7}{2}^+)$	$\Delta(\frac{9}{2}^+)$	$\Delta(\frac{11}{2}^+)$
		$n = 2$	~ 2390	(2390 ± 100)	(2360 ± 125)	(2460 ± 120)
		$n = 3$	~ 2610	~ 2610	~ 2610	~ 2610
		$n = 4$	~ 2820	~ 2820	~ 2820	~ 2820
$L = 6$	$S = \frac{1}{2}$	$n = 1$		$\Delta(\frac{11}{2}^+)$	$\Delta(\frac{13}{2}^+)$	
		$n = 2$		~ 2730	~ 2730	
		$n = 3$		~ 2930	~ 2930	
		$n = 4$				
	$S = \frac{3}{2}$	$n = 1$	$\Delta(\frac{9}{2}^+)$	$\Delta(\frac{11}{2}^+)$	$\Delta(\frac{13}{2}^+)$	$\Delta(\frac{15}{2}^+)$
		$n = 2$	~ 2810	~ 2810	~ 2810	(2920 ± 100)
		$n = 3$	~ 3000	~ 3000	~ 3000	~ 3000
		$n = 4$				

(33)

and for $P = -$ ones:

$L = 3$	$S = \frac{1}{2}$	$n = 1$	$\Delta(\frac{5}{2}^-)$	$\Delta(\frac{7}{2}^-)$		
		$n = 2$	~ 2230	(2230 ± 50)		
		$n = 3$	~ 2460	~ 2460		
		$n = 4$	~ 2670	~ 2670		
	$S = \frac{3}{2}$	$n = 1$	$\Delta(\frac{3}{2}^-)$	$\Delta(\frac{5}{2}^-)$	$\Delta(\frac{7}{2}^-)$	$\Delta(\frac{9}{2}^-)$
		$n = 2$	~ 2320	(2350 ± 50)	~ 2320	~ 2320
		$n = 3$	~ 2550	~ 2550	~ 2550	~ 2550
		$n = 4$	~ 2760	~ 2760	~ 2760	~ 2760
$L = 5$	$S = \frac{1}{2}$	$n = 1$		$\Delta(\frac{9}{2}^-)$	$\Delta(\frac{11}{2}^-)$	
		$n = 2$		(2400 ± 190)	~ 2670	
		$n = 3$		~ 2870	~ 2870	
		$n = 4$		~ 3050	~ 3050	
	$S = \frac{3}{2}$	$n = 1$	$\Delta(\frac{7}{2}^-)$	$\Delta(\frac{9}{2}^-)$	$\Delta(\frac{11}{2}^-)$	$\Delta(\frac{13}{2}^-)$
		$n = 2$	~ 2760	~ 2760	~ 2760	(2720 ± 100)
		$n = 3$	~ 2950	~ 2950	~ 2950	~ 2950
		$n = 4$	~ 3130	~ 3130	~ 3130	~ 3130

(34)

The above-written equations allow us to predict the (n, M^2) plots which are shown in Fig. 4.




Fig. 4. The (n, M^2) plots for $I = 1/2$ and $I = 3/2$ states.

6. Conclusion

Under the hypothesis of the quark–diquark structure of highly excited baryons, we succeeded to suggest for them a realistic classification. The introduction of diquarks gave a considerable reduction of excited states; additional reduction was obtained owing to the assumption of the overlapping of ($I = 1/2$) states with $S = 1/2$.

Thus obtained a classification gave linear trajectories in the (J, M^2) and (n, M^2) planes, which are strictly ordered. There is a number of overlapping poles. The observation of two-pole and three-pole structures in the complex- M planes of partial amplitudes is a top-priority task at the analysis of baryon spectra, while the next task consists in the writing and solving the spectral integral equation for quark–diquark systems. Such an equation should be similar to that written and solved before for the $q\bar{q}$ system¹⁹.

Acknowledgments

We thank L.G. Dakhno for helpful discussions. The paper was supported by the RFFI grants 07-02-01196-a and RSGSS-3628.2008.2.

References

1. N. Isgur and G. Karl, Phys. Rev. D **18**, 4187 (1978).
2. N. Isgur and G. Karl, Phys. Rev. D **19**, 2653 (1979).
3. S. Capstick, N. Isgur, Phys. Rev. D **34**, 2809 (1986).
4. L.Y. Glozman et al., Phys. Rev. D **58**:094030 (1998).
5. U. Löring, B.C. Metsch, H.R. Petry, Eur.Phys. A **10**, 395 (2001); A **10**, 447 (2001).
6. V.V. Anisovich, M.N. Kobrinsky, J. Nyiri, Yu.M. Shabelski *Quark Model and High Energy Collisions*, Second Edition, World Scientific, Singapore, 2004.
7. A.V. Anisovich, V.V. Anisovich, J. Nyiri, V.A. Nikonov, M.A. Matveev and A.V. Sarantsev, *Mesons and Baryons. Systematization and Methods of Analysis*, World Scientific, Singapore, 2008.
8. A.V. Anisovich, V.V. Anisovich, M.A. Matveev, V.A. Nikonov, A.V. Sarantsev and T.O. Vulfs *Searching for the quark-diquark systematics of baryons. Systems composed by the light quarks $q = u, d$* , arXiv:1001.1259 [hep-ph].
9. A.V. Sarantsev, *Methods of Data Analysis: Mesons and Baryons*, Talk given at the HADRON 2009, XIII Int. Conference on Hadron Spectroscopy, Tallahassee, 2009.
10. A.V. Anisovich, *Status of the Bonn-Gatchina partial wave analysis*, Talk given at the HADRON 2009, XIII Int. Conference on Hadron Spectroscopy, Tallahassee, 2009.
11. S.M. Flatté, Phys. Lett. B **63** 224 (1974).
12. C. Amsler et al. [Particle Data Group], Phys. Lett. B **667** (2008) 1.
13. R. A. Arndt, W. J. Briscoe, I. I. Strakovsky and R. L. Workman, Phys. Rev. C **74**, 045205 (2006) [arXiv:nucl-th/0605082].
14. R. E. Cutkosky, C. P. Forsyth, J. B. Babcock, R. L. Kelly and R. E. Hendrick,
15. D. M. Manley and E. M. Saleski, Phys. Rev. D **45**, 4002 (1992).
16. G. Hohler, In **Bratislava 1973, Proceedings, Triangle Meeting On Hadron Interactions At Low Energies**, Bratislava 1975, 11-75 and Preprint - HOEHLER G (73,REC.DEC) 60p
17. A.V. Anisovich, V.V. Anisovich, and A.V. Sarantsev, Phys. Rev. D **62**:051502(R) (2000).

18. E. Klempt, *Baryon Spectroscopy and the Origin of Mass*, Talk given at the HADRON 2009, XIII Int. Conference on Hadron Spectroscopy, Tallahassee, 2009.
19. V.V. Anisovich, L.G. Dakhno, M.A. Matveev, V.A. Nikonov, and A. V. Sarantsev, *Yad. Fiz.* **70**, 480 (2007) [*Phys. Atom. Nucl.* **70**, 450 (2007)].